

Trying 3106016892...Open

Welcome to STN International! Enter x:x  
LOGINID:ssspta1205mxm  
PASSWORD:  
TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Feb 2 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Dec 17 Expanded CPlus Coverage of US, Japanese, WIPO,  
EPO, and German patents  
NEWS 3 Feb 1 Addition of Machine-Translated Abstracts to CPlus  
NEWS 4 Feb 2 STEREO BOND SEARCH PROBLEM FIXED WITH STN EXPRESS 5.0C  
NEWS 5 Feb 14 Homology Searching for Nucleotide Sequences in DGENE  
now available!  
NEWS 6 Feb 16 BIOTECHNOBASE NOW ON STN  
NEWS 7 Feb 22 New Database Producer Clusters Now Available on STN  
NEWS 8 Feb 28 Structure Search Limits Increased in REGISTRY,  
ZREGISTRY, and CASREACT  
NEWS 9 Feb 28 Patent Information Now Searchable in CAOLD  
NEWS 10 Mar 1 New IMSDIRECTORY Provides Pharma Company Details  
NEWS 11 Mar 20 INPADOC: PRODUCER WARNING ABOUT DATA DELAYS  
NEWS 12 Mar 22 NEW FEATURES IN INPADOC - RANGE SEARCHING AND NEW  
SDI/UPDATE SEARCH FIELD

NEWS EXPRESS FREE UPGRADE 5.0C NOW AVAILABLE  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 14:40:01 ON 26 MAR 2000

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.15	0.15

FILE 'REGISTRY' ENTERED AT 14:40:07 ON 26 MAR 2000  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2000 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 24 MAR 2000 HIGHEST RN 259886-75-4

prov.  
2/23/98

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 13, 1999

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

Structure search limits have been increased. See HELP SLIMIT  
for details.

=> s viagra/cn

L1 1 VIAGRA/CN

=> s sildenafil?/cn

L2 2 SILDENAFIL?/CN

=> file medline, hcaplus, uspatfull

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	7.50	7.65

FILE 'MEDLINE' ENTERED AT 14:40:46 ON 26 MAR 2000

FILE 'HCAPLUS' ENTERED AT 14:40:46 ON 26 MAR 2000  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2000 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'USPATFULL' ENTERED AT 14:40:46 ON 26 MAR 2000  
CA INDEXING COPYRIGHT (C) 2000 AMERICAN CHEMICAL SOCIETY (ACS)

=> s l1 or l2

L3 445 L1 OR L2

=> s ed or ereecti? or impoten?

L4 228938 ED OR EERECTI? OR IMPOTEN?

=> s ed or erecti? or impoten?

L5 246668 ED OR ERECTI? OR IMPOTEN?

=> s spine or spinal or paraly?

L6 267236 SPINE OR SPINAL OR PARALY?

=> s l3 and l5

L7 345 L3 AND L5

=> s l7 and l6

L8 16 L7 AND L6

=> dup rem l8

PROCESSING COMPLETED FOR L8

L9 12 DUP REM L8 (4 DUPLICATES REMOVED)

=> d bib,ab,kwic l9 1-12

L9 ANSWER 1 OF 12 USPATFULL  
AN 2000:31420 USPATFULL  
TI Local administration of phosphodiesterase inhibitors for the treatment  
of **erectile dysfunction**  
IN Doherty, Jr., Paul C., Cupertino, CA, United States  
Place, Virgil A., Kawaihae, HI, United States  
Smith, William L., Mahwah, NJ, United States  
PA Vivus, Inc., Mountain View, CA, United States (U.S. corporation)  
PI US 6037346 20000314  
AI US 1998-181070 19981027 (9)  
RLI Continuation-in-part of Ser. No. US 1997-958816, filed on 28 Oct 1997,  
now abandoned  
DT Utility  
EXNAM Primary Examiner: Reamer, James H.  
LREP Reed, Dianne E. Reed & Associates  
CLMN Number of Claims: 94  
ECL Exemplary Claim: 1,23  
DRWN 1 Drawing Figure(s); 1 Drawing Page(s)  
LN.CNT 1331

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB A method is provided for treating **erectile dysfunction** in a  
mammalian male individual. The method involves the local administration  
of a phosphodiesterase inhibitor or a pharmaceutically acceptable salt,  
ester, amide or derivative thereof within the context of an effective  
dosing regimen. A preferred mode of administration is transurethral.  
Pharmaceutical formulations and kits are provided as well.  
TI Local administration of phosphodiesterase inhibitors for the treatment  
of **erectile dysfunction**  
AB A method is provided for treating **erectile dysfunction** in a  
mammalian male individual. The method involves the local administration  
of a phosphodiesterase inhibitor or a pharmaceutically acceptable.

SUMM This invention relates generally to methods and pharmaceutical  
compositions for treating **erectile dysfunction**; more  
particularly, the invention relates to local administration of  
phosphodiesterase inhibitors to treat **erectile dysfunction**.

SUMM **Impotence** is the consistent inability to achieve or sustain an  
**erection** of sufficient rigidity for sexual intercourse. It has  
recently been estimated that approximately 10 million American men are  
**impotent** (R. Shabsigh et al., "Evaluation of **Erectile**  
**Impotence**," Urology 32:83-90 (1988); W. L. Furlow, "Prevalence  
of **Impotence** in the United States," Med Aspects Hum. Sex.  
19:13-6 (1985)). **Impotence** is recognized to be an  
age-dependent disorder, with an incidence of 1.9 percent at 40 years of  
age and 25. . . Male; A. C. Kinsey et al., eds., Philadelphia, Pa.:  
W. B. Saunders, 218-262 (1948)). In 1985 in the United States,  
**impotence** accounted for more than several hundred thousand  
outpatient visits to physicians Rational Center for Health Statistics,  
National Hospital Discharge Survey, . . .

SUMM A number of causes of **impotence** have been identified,  
including vasculogenic, neurogenic, endocrinologic and psychogenic.  
Vasculogenic **impotence**, which is caused by alterations in the  
flow of blood to and from the penis, is thought to be the most frequent  
organic cause of **impotence**. Common risk factors for  
vasculogenic **impotence** include hypertension, diabetes,  
cigarette smoking, pelvic trauma, and the like. Neurogenic  
**impotence** is associated with spinal-cord injury,  
multiple sclerosis, peripheral neuropathy caused by diabetes or  
alcoholism and severance of the autonomic nerve supply to the penis  
consequent to prostate surgery. **Erectile dysfunction** is also  
associated with disturbances in endocrine function resulting in low  
circulating testosterone levels and elevated prolactin levels.

SUMM **Impotence** can also be a side effect of various classes of  
drugs, in particular, those that interfere with central neuroendocrine

control. . . or local neurovascular control of penile smooth muscle. Krane et al., New England Journal of Medicine 321: 1648 (1989). Penile **erection** requires (1) dilation of the arteries that regulate blood flow to the lacunae of the corpora cavernosum, (2) relaxation of.

SUMM . . . vasoactive substances such as vasoactive intestinal polypeptide (VIP), prostanoids, endothelin and nitric oxide. High sympathetic tone (noradrenergic) is implicated in **erectile** dysfunction, and, in some patients, the disorder can be successfully treated with noradrenergic receptor antagonists. See, e.g., Krane et al., . . .

SUMM There is also evidence that dopaminergic mechanisms are involved in **erectile** function. For example, pharmacologic agents that elevate the level of brain dopamine or stimulate brain dopamine receptors increase sexual activity. . . .

SUMM . . . Hyppa et al., Acta Neurologic Scand. 46:223 (Supp. 43, 1970)). Specific dopamine agonists have been studied for their effects on **erectile** function. Apomorphine, (n-propyl)norapo-morphine, bromocryptine, amantidine, fenfluramine, L-DOPA and various other pharmacological activators of central dopaminergic receptors have been found to increase episodes of penile **erection** in male rats (Benassi-Benelli et al., Arch. int. Pharmacodyn. 242:241 (1979); Poggioli et al., Riv. di Farm. & Terap. 9:213. . . .

SUMM The currently available dopamine agonists, with few exceptions, have found limited use in the treatment of **erectile** dysfunction because of their peripheral side effects. These effects include nausea and vomiting, postural hypotension, arrhythmias, tachycardia, dysphoria, psychosis, hallucinations, drowsiness and dysidnesias (See, e.g., Martindale The Extra Pharmacopoeia, 31st Ed., pages 1151-1168).

SUMM Other pharmaceutical methods for treating **erectile** dysfunction have also proved to be problematic. For example, with Viagra.RTM., the most recently introduced oral drug therapy, not only. . . .

SUMM . . . herein provides a means to avoid the above-mentioned problems encountered with the systemic administration of pharmacologically active agents to treat **erectile** dysfunction. Specifically, the invention relates to methods and formulations for effectively treating **erectile** dysfunction by locally administering a selected active agent, wherein the active agent is an inhibitor of a phosphodiesterase.

SUMM . . . messenger nucleotides, cyclic adenosine monophosphate (cAMP), and cyclic guanosine monophosphate (cGMP) (see, e.g., Doherty, "Oral, Transdermal, and Transurethral Therapies for **Erectile** Dysfunction" in Male Infertility and Dysfunction, Hellstrom, ed ., Chapter 34 (New York, N.Y.: Springer-VerlagHellstrom, 1997)). Numerous phosphodiesterase inhibitors have previously been described in the literature for a variety. . . . 34). Oral and parenteral administration of phosphodiesterase inhibitors, as alluded to above, have also been suggested for the treatment of **erectile** dysfunction (Doherty, supra; see also PCT Publication Nos. WO 96/16644, and WO 94/28902). The phosphodiesterases have been classified into seven. . . .

SUMM The following documents are of interest insofar as they relate to the treatment of **erectile** dysfunction by delivering pharmacologically active agents locally to the penis:

SUMM . . . injection of vasodilator drugs into the corpora cavernosa of the penis to dilate the arteries that supply blood to the **erectile** tissues, thereby inducing an **erection**;

SUMM . . . direct injection of a drug into the corpora cavernosa, by topical drug administration or transurethral drug administration, for inhibiting penile **erection** due to priapism and for treating urinary incontinence;

SUMM . . . the intracavernosal injection of papaverine (a smooth muscle relaxant), phenoxybenzamine or phentolamine (.alpha.-receptor blockers)

and a phentolamine-papaverine mixture to treat **erectile** dysfunction; and

SUMM . . . et al., and U.S. Pat. Nos. 5,242,391, 5,474,535, 5,686,093 and 5,773,020 to Place et al. relate to the treatment of **erectile** dysfunction by delivery of a vasoactive agent into the male urethra.

SUMM The invention, as noted above, is directed to local administration of pharmacologically active agents to treat **erectile** dysfunction. The agents are preferably, although not necessarily, Type V phosphodiesterase inhibitors. Surprisingly, it has now been found by

the inventors herein that local administration of these phosphodiesterase inhibitors as disclosed herein is highly effective in treating **erectile** dysfunction, particularly vasculogenic **impotence**. Local administration of phosphodiesterase inhibitors, and transurethral drug administration in particular, generally enables use of a lower drug dosage, avoids . . . administered medications an individual may be taking. The local administration of phosphodiesterase inhibitors, particularly Type V phosphodiesterase inhibitors, to treat **erectile** dysfunction, accordingly represents an important advance in the treatment of **impotence** and other **erectile** disorders.

SUMM . . . primary object of the invention to address the above-described need in the art by providing a novel method for treating **erectile** dysfunction by locally administering an effective amount of a selected phosphodiesterase inhibitor to an individual in need of such therapy.

SUMM In a first aspect of the invention, a method is provided for treating an individual prone to **erectile** dysfunction, particularly vasculogenic **erectile** dysfunction, the method comprising locally administering to the individual a pharmaceutical formulation containing a phosphodiesterase inhibitor. Administration of the pharmaceutical . . . carried out within the context of a predetermined dosing regimen such that the agent is effective in the treatment of **erectile** dysfunction. The method is especially useful in the treatment of vasculogenic **impotence**, although other types of **erectile** dysfunction may also be treated using the present formulations. Drug delivery is preferably effected transurethally, but the drug may also. . .

SUMM In another aspect of the invention, a pharmaceutical formulation is provided for carrying out the present method for treating **erectile** dysfunction. The pharmaceutical formulation comprises an effective amount of a phosphodiesterase inhibitor, a carrier or vehicle preferably suitable for the. . .

SUMM . . . formulation during storage and prior to use; and instructions for carrying out drug administration in a manner effective to treat **erectile** dysfunction.

DETD The term "**erectile** dysfunction" is intended to include any and all types of **erectile** dysfunction, including: vasculogenic, neurogenic, endocrinologic and psychogenic **impotence** ("**impotence**" is used here in its broadest sense to indicate an inability a periodic or consistent inability to achieve or sustain an **erection** of sufficient rigidity for sexual intercourse; see U.S. Pat. No. 5,242,391 to Place et al., cited supra); Peyronie's syndrome; priapism; . . .

DETD . . . of the occurrence of symptoms and/or their underlying cause, and improvement or remediation of damage. The present method of "treating" **erectile** dysfunction, as the term is used herein, thus encompasses both prevention of the disorder in a predisposed individual and treatment. . .

DETD . . . is meant a nontoxic but sufficient amount of the drug or agent to provide the desired effect, i.e., treatment of **erectile** dysfunction.

DETD Active Agents for Treatment of **Erectile** Dysfunction:

DETD . . . to carry out the method of the invention, a selected phosphodiesterase inhibitor is locally administered to an individual prone to **erectile** dysfunction.

DETD . . . art of synthetic organic chemistry and described, for example, by J. March, Advanced Organic Chemistry; Reactions, Mechanisms and Structure, 4th Ed. (New York: Wiley-Interscience, 1992). For example, acid addition salts are prepared from the free base using conventional methodology, and involves. . .

DETD The active agent is administered locally to treat **erectile** dysfunction, and is accordingly administered in a pharmaceutical formulation suitable for local administration.

DETD . . . as described in the pertinent literature and pharmaceutical texts. See, for example, Remington: The Science and Practice of Pharmacy, 19th Ed. (Easton, Pa.: Mack Publishing Co., 1995), which discloses typical methods of preparing pharmaceutical compositions

in the form of urethral suppositories.. . .

DETD . . . control device such as that described in PCT Publication No. WO

97/47260, entitled "Venous Flow Control Element for Maintaining Penile **Erection**." Preferred devices are formed from a length of flexible tubing having an integral fastening means, so as to provide for. . . it effectively enhances retention of blood within the penis without substantially obstructing arterial inflow or becoming too constrictive during the **erectile** process. Use of the VFC device also enables enhanced effectiveness of local drug therapy, in that the active agent is retained within the penis, allowing movement into the corpus cavernosa. This produces smooth muscle response and a consistent **erectile** response. In this embodiment, a kit will include the venous flow control device in addition to the components noted above,. . .

DETD The pharmaceutical preparations of Examples 1 and 2 can be used to treat **erectile** dysfunction in individuals in which the dysfunction is associated, for example, vascular insufficiency. Dosage may be adjusted using the methodology. . . Example 3. In all instances the individuals are expected to respond positively, although variations in the intensity and duration of **erection** may be observed depending on dose, formulation and environment. Generally, between approximately 20 and 90 minutes following drug administration, it is expected that an **erection** may be achieved.

DETD In this experiment, zaprinast, a Type V phosphodiesterase inhibitor, was evaluated for its capability to induce **erections** in the anesthetized male cat. Adult male cats (3.5 to 5.0 kg) were initially sedated with ketamine and then anesthetized. . . mm, respectively). These results suggest that a selective Type V phosphodiesterase inhibitor, when administered locally, can induce significant increases in **erectile** response in a mammalian male. The same or greater effects are expected upon administration of a urethral suppository.

CLM What is claimed is:

1. A method for treating **erectile** dysfunction in a male individual, comprising locally administering to the individual an effective amount of a pharmaceutical composition consisting essentially.
13. The method of claim 1, wherein the **erectile** dysfunction is vasculogenic **impotence**.
18. A pharmaceutical formulation for treating **erectile** dysfunction in an individual, comprising a urethral dosage form of a phosphodiesterase inhibitor, a carrier suitable for transurethral drug administration,. . .
20. A pharmaceutical formulation for treating **erectile** dysfunction in an individual comprising a sterile liquid composition

suitable for intracavernosal administration containing a therapeutically

effective amount of a. . .

21. A pharmaceutical formulation for treating **erectile** dysfunction in an individual, comprising a topical or transdermal composition containing a therapeutically effective amount of a phosphodiesterase inhibitor and. . .

23. A kit for treating **erectile** dysfunction in an individual, comprising: a urethral dosage form of a Type V, cGMP-specific phosphodiesterase inhibitor or a pharmaceutically acceptable. . .

for

using the drug delivery means to administer the drug within the context of a dosing regimen effective to treat **erectile** dysfunction.

86. A method for treating **erectile** dysfunction in a male individual, comprising administering to the individual a pharmaceutical composition comprising a therapeutically effective amount of: (a). .

IT 58-32-2, Dipyridamole 69-89-6D, Xanthine, derivs. 120-73-0D, Purine, derivs. 253-82-7D, Quinazoline, derivs. 289-95-2D, Pyrimidine, derivs. 37762-06-4, Zaprinast 51022-77-6, Etazolate 51022-77-6D, Etazolate, esters and analogs 56739-21-0, Nitraquazone 56739-21-0D, Nitraquazone, esters and analogs 57076-71-8, Denbufylline  
61413-54-5,  
Rolipram 61413-54-5D, Rolipram, esters and analogs 66327-51-3, Furazlocillin 79030-08-3D, Griseolic acid, derivs. 120223-30-5, EMD54622 120223-30-5D, EMD54622, esters and analogs 136145-07-8, LAS-31025 136145-07-8D, LAS-31025, esters and analogs 139145-27-0  
139755-83-2, Sildenafil 147676-63-9 150452-19-0 167298-74-0  
184147-55-5 190281-17-5D, Pyrazolopyrimidinone, derivs. 224157-99-7  
(phosphodiesterase inhibitor local administration for treatment of **erectile** dysfunction)

L9 ANSWER 2 OF 12 MEDLINE

AN 1999334975 MEDLINE

DN 99334975

TI **Erectile** dysfunction in spina bifida is treatable [letter].

AU Palmer J S; Kaplan W E; Firlit C F

SO LANCET, (1999 Jul 10) 354 (9173) 125-6.

Journal code: LOS. ISSN: 0140-6736.

CY ENGLAND: United Kingdom

DT (CLINICAL TRIAL)

Letter

(RANDOMIZED CONTROLLED TRIAL)

LA English

FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals

EM 199910

EW 19991001

AB We undertook a prospective, blinded, randomised, placebo-controlled, dose escalation, crossover study that showed that **erectile** dysfunction in spina bifida is medically treatable, specifically with sildenafil citrate.

TI **Erectile** dysfunction in spina bifida is treatable [letter].

AB We undertook a prospective, blinded, randomised, placebo-controlled, dose escalation, crossover study that showed that **erectile** dysfunction in spina bifida is medically treatable, specifically with sildenafil citrate.

CT Check Tags: Human; Support, Non-U.S. Gov't

Adult

Analysis of Variance

Cross-Over Studies

\*Impotence: DT, drug therapy

Impotence: ET, etiology

\*Phosphodiesterase Inhibitors: TU, therapeutic use

\*Piperazines: TU, therapeutic use

Single-Blind Method  
\*Spinal Dysraphism: CO, complications  
RN 139755-83-2 (sildenafil)

L9 ANSWER 3 OF 12 HCAPLUS COPYRIGHT 2000 ACS  
AN 1999:690785 HCAPLUS  
DN 131:281606  
TI Method of treating **impotence** due to **spinal** cord injury  
with sildenafil or other cGMP phosphodiesterase inhibitor  
IN Maytom, Murray Craig; Osterloh, Ian Howard  
PA Pfizer Ltd., UK; Pfizer Research and Development Company, N.V./S.A.  
SO Eur. Pat. Appl., 7 pp.  
CODEN: EPXXDW  
DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 951908	A2	19991027	EP 1999-301085	19990215
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	JP 11315025	A2	19991116	JP 1999-43205	19990222
	AU 9918390	A1	19990909	AU 1999-18390	19990223
PRAI	US 1998-75580		19980223		

OS MARPAT 131:281606

AB A class of cGMP phosphodiesterase inhibitors, including sildenafil and pharmaceutically acceptable salts thereof, is disclosed for use in the treatment of sexual dysfunction in male and female animals, esp. humans, with a **spinal** cord injury. The invention can be used to treat sexual dysfunction in male animals that exhibit essentially no residual penile function.

TI Method of treating **impotence** due to **spinal** cord injury with sildenafil or other cGMP phosphodiesterase inhibitor

AB . . . thereof, is disclosed for use in the treatment of sexual dysfunction in male and female animals, esp. humans, with a **spinal** cord injury. The invention can be used to treat sexual dysfunction in male animals that exhibit essentially no residual penile. . . .

ST cGMP phosphodiesterase inhibitor sexual dysfunction; sildenafil sexual dysfunction **spinal** cord injury; **erectile** dysfunction  
**spinal** cord injury sildenafil

IT Sexual behavior  
(disorder; sildenafil or other cGMP phosphodiesterase inhibitor for treatment of sexual dysfunction in animal with **spinal** cord injury)

IT Sexual behavior  
(**impotence**; sildenafil or other cGMP phosphodiesterase inhibitor for treatment of sexual dysfunction in animal with **spinal** cord injury)

IT **Spinal** cord  
(injury; sildenafil or other cGMP phosphodiesterase inhibitor for treatment of sexual dysfunction in animal with **spinal** cord injury)

IT 9068-52-4, CGMP phosphodiesterase

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(inhibitors; sildenafil or other cGMP phosphodiesterase inhibitor for treatment of sexual dysfunction in animal with **spinal** cord injury)

IT 139755-83-2 171599-83-0, Sildenafil citrate

RL: BAC (Biological activity or effector, except adverse); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(sildenafil or other cGMP phosphodiesterase inhibitor for treatment of sexual dysfunction in animal with **spinal** cord injury)

L9 ANSWER 4 OF 12 MEDLINE  
AN 1999325995 MEDLINE

DUPLICATE 1



DN 99325995  
TI Sildenafil: a review of its use in **erectile** dysfunction.  
AU Langtry H D; Markham A  
CS Adis International Limited, Mairangi Bay, Auckland, New Zealand.  
SO DRUGS, (1999 Jun) 57 (6) 967-89. Ref: 80  
Journal code: EC2. ISSN: 0012-6667.  
CY New Zealand  
DT Journal; Article; (JOURNAL ARTICLE)  
General Review; (REVIEW)  
(REVIEW, ACADEMIC)  
LA English  
FS Priority Journals  
EM 199911  
EW 19991102  
AB Sildenafil is an oral therapy for **erectile** dysfunction of a  
broad range of causes. By selectively inhibiting phosphodiesterase type  
5,

it allows corpus cavernosum smooth muscle to relax, potentiating  
**erections** during sexual stimulation. Blood pressure is reduced  
transiently by sildenafil, but more marked hypotension may occur during  
concurrent administration of sildenafil and organic nitrates; this  
combination is contraindicated. Sildenafil is rapidly absorbed, with  
dose-proportional peak plasma concentrations within 1 hour of  
administration. The elimination half-life is 3 to 5 hours. Dosages  
usually  
begin at 50mg taken when needed =1 hour before sexual activity no more  
than once daily. The maximum dose is 100mg when needed once daily and  
lower doses (e.g. 25mg) may be used in elderly patients and those with  
hepatic or renal impairment or receiving cytochrome P450 enzyme CYP3A4  
inhibitors, such as ritonavir, saquinavir, ketoconazole, erythromycin or  
cimetidine. More than 3000 patients with **erectile** dysfunction of  
organic (e.g. diabetes or **spinal** cord injury), psychogenic or  
mixed origin received sildenafil 5 to 100mg or placebo in fixed- or  
titrated-dose trials. Sildenafil was associated with dose-related  
improvements in the frequency, hardness and duration of **erections**  
and in patients' abilities to achieve and maintain **erections**  
adequate for successful sexual intercourse. In titrated-dose trials, the  
most commonly effective doses were 50 or 100mg, although lower doses were  
effective in some patients. Sildenafil was significantly more effective  
than placebo in **erectile** dysfunction of all tested causes. The  
efficacy of sildenafil was not affected by patient age (> or < or =65  
years) or by antihypertensive or antidepressant medications. The drug was  
effective in patients with severe **erectile** dysfunction. Efficacy  
was maintained in long term (1-year) studies. Sildenafil also appears to  
improve the quality of life of both patients and their sexual partners.  
Common adverse events associated with sildenafil were transient and mild  
or moderate and included headache, flushing, dyspepsia, nasal congestion  
and abnormal vision. Tolerability was maintained in long term (< or =1  
year) studies. No serious sildenafil-related adverse events occurred in  
clinical trials; cardiovascular events seen in postmarketing surveillance  
generally occurred in patients with other known risk factors.

#### CONCLUSIONS:

Sildenafil is an effective oral treatment in men with **erectile**  
dysfunction. It was significantly superior to placebo in improving  
**erections** and allowing successful penetrative sexual intercourse.  
Although its place in disease management is still emerging and there are  
contraindications to its use, if preliminary positive reports are  
confirmed, sildenafil will be the pre-eminent first-line therapy for  
**erectile** dysfunction.

TI Sildenafil: a review of its use in **erectile** dysfunction.  
AB Sildenafil is an oral therapy for **erectile** dysfunction of a  
broad range of causes. By selectively inhibiting phosphodiesterase type  
5,  
it allows corpus cavernosum smooth muscle to relax, potentiating  
**erections** during sexual stimulation. Blood pressure is reduced

transiently by sildenafil, but more marked hypotension may occur during concurrent administration of. . . or receiving cytochrome P450 enzyme CYP3A4 inhibitors, such as ritonavir, saquinavir, ketoconazole, erythromycin or cimetidine. More than 3000 patients with **erectile** dysfunction of organic (e.g. diabetes or **spinal** cord injury), psychogenic or mixed origin received sildenafil 5 to 100mg or placebo in fixed- or titrated-dose trials. Sildenafil was associated with dose-related improvements in the frequency, hardness and duration of **erections** and in patients' abilities to achieve and maintain **erections** adequate for successful sexual intercourse. In titrated-dose trials, the most commonly effective doses were 50 or 100mg, although lower doses were effective in some patients. Sildenafil was significantly more effective than placebo in **erectile** dysfunction of all tested causes. The efficacy of sildenafil was not affected by patient age (> or < or =65 years) or by antihypertensive or antidepressant medications. The drug was effective in patients with severe

**erectile** dysfunction. Efficacy was maintained in long term (1-year) studies. Sildenafil also appears to improve the quality of life of both. . . surveillance generally occurred in patients with other known risk factors. CONCLUSIONS: Sildenafil is an effective oral treatment

in men with **erectile** dysfunction. It was significantly superior to placebo in improving **erections** and allowing successful penetrative sexual intercourse. Although its place in disease management is still emerging and there are contraindications to its use, if preliminary positive reports are confirmed, sildenafil will be the pre-eminent first-line therapy for **erectile** dysfunction.

CT Check Tags: Human; Male  
Clinical Trials  
Drug Interactions  
\*Impotence: DT, drug therapy  
Phosphodiesterase Inhibitors: AE, adverse effects  
Phosphodiesterase Inhibitors: PD, pharmacology  
\*Phosphodiesterase Inhibitors: TU, therapeutic use  
\*Piperazines: TU, therapeutic. . .

RN 139755-83-2 (sildenafil)

L9 ANSWER 5 OF 12 HCAPLUS COPYRIGHT 2000 ACS  
AN 2000:15560 HCAPLUS  
DN 132:59075  
TI Sildenafil citrate to treat **erectile** dysfunction in the spina bifida male  
AU Palmer, Jeffrey S.; Kaplan, William E.; Firlit, Casimir F.  
CS Division of Urology, Children's Memorial Medical Center, Northwestern University Medical School, Chicago, IL, USA  
SO Surg. Forum (1999), 50, 712-713  
CODEN: SUFOAX; ISSN: 0071-8041  
PB American College of Surgeons  
DT Journal  
LA English  
AB Sildenafil citrate (25 or 50 mg) dose-dependently improved **erectile** function in men with spina bifida.  
TI Sildenafil citrate to treat **erectile** dysfunction in the spina bifida male  
AB Sildenafil citrate (25 or 50 mg) dose-dependently improved **erectile** function in men with spina bifida.  
ST sildenafil **erectile** dysfunction spina bifida; **impotence** spina bifida sildenafil  
IT Sexual behavior  
(**impotence**; sildenafil citrate treatment of **erectile** dysfunction in men with spina bifida)  
IT **Spinal** column  
(spina bifida; sildenafil citrate treatment of **erectile** dysfunction in men with spina bifida)

IT 171599-83-0, Sildenafil citrate  
 RL: BAC (Biological activity or effector, except adverse); THU  
 (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (sildenafil citrate treatment of **erectile** dysfunction in men  
 with spina bifida)

L9 ANSWER 6 OF 12 MEDLINE  
 AN 1999163632 MEDLINE  
 DN 99163632  
 TI A two-part pilot study of sildenafil (VIAGRA) in men with **erectile**  
 dysfunction caused by **spinal** cord injury.  
 AU Maytom M C; Derry F A; Dinsmore W W; Glass C A; Smith M D; Orr M;  
 Osterloh  
 I H  
 CS Pfizer Central Research, Sandwich, UK.  
 SO SPINAL CORD, (1999 Feb) 37 (2) 110-6.  
 Journal code: CKK. ISSN: 1362-4393.  
 CY ENGLAND: United Kingdom  
 DT (CLINICAL TRIAL)  
 Journal; Article; (JOURNAL ARTICLE)  
 (MULTICENTER STUDY)  
 (RANDOMIZED CONTROLLED TRIAL)  
 LA English  
 FS Priority Journals  
 EM 199907  
 EW 19990702  
 AB STUDY DESIGN: This was a two-part pilot study in men with **erectile**  
 dysfunction (**ED**) due to **spinal** cord injury (SCI: cord  
 level range T6-L5). Part I was a randomised, double-blind, two-way  
 cross-over study comparing a single dose of sildenafil 50 mg or placebo.  
 Part II was a randomised, double-blind, parallel-group evaluation of  
 sildenafil 50 mg or placebo, taken as required (not more than once daily)  
 approximately 1 h prior to sexual activity, over a period of 28 days.  
 OBJECTIVES: To assay the efficacy and safety of sildenafil 50 mg and  
 placebo. SETTING: Clinic- and home-based assessments in the United  
 Kingdom. METHODS: A total of 27 subjects who were able to achieve at  
 least  
 a grade 2 **erection** (hard, but not hard enough for penetration)  
 in response to penile vibratory stimulation (PVS) were recruited. In Part  
 I, the reflexogenic response of the penis to PVS was evaluated in the  
 clinic while in Part II, the response to treatment was assessed in the  
 home (global efficacy. questionnaire, diary). RESULTS: In Part I, 17/26  
 (65%) subjects had **erections** of >60% rigidity at the penile base  
 (median duration 3.5 min) after sildenafil compared with 2/26 (8%)  
 (median  
 duration 0 min) after placebo (P=0.0003). In Part II, 9/12 (75%) subjects  
 on sildenafil and 1/14 (7%) subjects on placebo reported that the  
 treatment had improved their **erections** (P<0.005), and 8/12 (67%)  
 and 2/13 (15%) men, respectively, indicated that they wished to continue  
 treatment (P<0.02). An analysis of diary data showed no difference  
 between  
 the groups with respect to the mean number of **erections** hard  
 enough for penetration (P = 0.08). The mean proportion of attempts at  
 sexual intercourse that were successful was 30 and 15%, respectively  
 (P=0.21). Similarly, responses to the end-of-treatment questionnaire  
 indicated that there were no significant differences between the groups  
 with respect to the frequency of **erections** hard enough for  
 sexual intercourse (P=0.47) or that lasted as long as the subject would  
 have liked (P=0.11). No subject discontinued sildenafil due to adverse  
 events. CONCLUSION: Sildenafil is an effective, well-tolerated oral  
 treatment for **ED** in SCI subjects.  
 TI A two-part pilot study of sildenafil (VIAGRA) in men with **erectile**  
 dysfunction caused by **spinal** cord injury.  
 AB STUDY DESIGN: This was a two-part pilot study in men with **erectile**  
 dysfunction (**ED**) due to **spinal** cord injury (SCI: cord

level range T6-L5). Part I was a randomised, double-blind, two-way cross-over study comparing a single dose. . . in the United Kingdom. METHODS: A total of 27 subjects who were able to achieve at least a grade 2 **erection** (hard, but not hard enough for penetration) in response to penile vibratory stimulation (PVS) were recruited. In Part I, the. . . response to treatment was assessed in the home (global efficacy. questionnaire, diary). RESULTS: In Part I, 17/26 (65%) subjects had **erections** of >60% rigidity at the penile base (median duration 3.5 min) after sildenafil compared with 2/26 (8%) (median duration 0. . . Part II, 9/12 (75%) subjects on sildenafil and 1/14 (7%) subjects on placebo reported that the treatment had improved their **erections** ( $P<0.005$ ), and 8/12 (67%) and 2/13 (15%) men, respectively, indicated that they wished to continue treatment ( $P<0.02$ ). An analysis of diary data showed no difference between the groups with respect to the mean number of **erections** hard enough for penetration ( $P = 0.08$ ). The mean proportion of attempts at sexual intercourse that were successful was 30. . . to the end-of-treatment questionnaire indicated that there were no significant differences between

the groups with respect to the frequency of **erections** hard enough for sexual intercourse ( $P=0.47$ ) or that lasted as long as the subject would have liked ( $P=0.11$ ). No subject discontinued sildenafil due to adverse events. CONCLUSION: Sildenafil is an effective, well-tolerated oral treatment for **ED** in SCI subjects.

CT Check Tags: Human; Male; Support, Non-U.S. Gov't  
Adult  
Cross-Over Studies  
Double-Blind Method  
\***Impotence: DT, drug therapy**  
\***Impotence: ET, etiology**  
Middle Age  
Phosphodiesterase Inhibitors: AE, adverse effects  
\*Phosphodiesterase Inhibitors: TU, therapeutic use  
Pilot Projects  
Piperazines: AE, adverse effects  
\*Piperazines: TU, therapeutic use  
\***Spinal Cord Injuries: CO, complications**  
Treatment Outcome

RN 139755-83-2 (sildenafil)

L9 ANSWER 7 OF 12 MEDLINE  
AN 2000128503 MEDLINE  
DN 20128503  
TI Sildenafil citrate (VIAGRA): a novel oral treatment for **erectile** dysfunction caused by traumatic **spinal** cord injury.  
AU Giuliano F; Hultling C; el Masry W S; Luchner E; Stien R; Maytom M C; Orr M; Smith M D; Osterloh I H  
CS Hopital de Bicetre, Paris, France.  
SO INTERNATIONAL JOURNAL OF CLINICAL PRACTICE. SUPPLEMENT, (1999 Jun) 102 24-6.  
Journal code: CW2. ISSN: 1368-504X.  
CY ENGLAND: United Kingdom  
DT (CLINICAL TRIAL)  
Journal; Article; (JOURNAL ARTICLE)  
(RANDOMIZED CONTROLLED TRIAL)  
LA English  
EM 200004  
EW 20000404  
TI Sildenafil citrate (VIAGRA): a novel oral treatment for **erectile** dysfunction caused by traumatic **spinal** cord injury.  
CT Check Tags: Human; Male  
Administration, Oral  
Adult  
Cross-Over Studies  
Double-Blind Method

\*Impotence: DT, drug therapy  
 Impotence: ET, etiology  
 Middle Age  
 \*Phosphodiesterase Inhibitors: AD, administration & dosage  
 \*Piperazines: AD, administration & dosage  
 Spinal Cord Injuries: CO, complications  
 RN 139755-83-2 (sildenafil)

L9 ANSWER 8 OF 12 MEDLINE  
 AN 2000128500 MEDLINE  
 DN 20128500  
 TI Partners' perceptions of the efficacy of sildenafil citrate (VIAGRA) in the treatment of **erectile** dysfunction.  
 AU Hultling C  
 CS Spinalis SCI Research Unit, Karolinska Institute, Stockholm, Sweden.  
 SO INTERNATIONAL JOURNAL OF CLINICAL PRACTICE. SUPPLEMENT, (1999 Jun) 102 16-8.  
 Journal code: CW2. ISSN: 1368-504X.  
 CY ENGLAND: United Kingdom  
 DT (CLINICAL TRIAL)  
 Journal; Article; (JOURNAL ARTICLE)  
 (RANDOMIZED CONTROLLED TRIAL)  
 LA English  
 EM 200004  
 EW 20000404  
 TI Partners' perceptions of the efficacy of sildenafil citrate (VIAGRA) in the treatment of **erectile** dysfunction.  
 CT Check Tags: Female; Human; Male  
 Adult  
 Aged  
 Aged, 80 and over  
 Cross-Over Studies  
 Double-Blind Method  
 \*Impotence: DT, drug therapy  
 Impotence: ET, etiology  
 Middle Age  
 \*Penile Erection  
 Perception  
 \*Phosphodiesterase Inhibitors: AD, administration & dosage  
 \*Piperazines: AD, administration & dosage  
 \*Sexual Partners  
 Spinal Cord Injuries: CO, complications  
 RN 139755-83-2 (sildenafil)

L9 ANSWER 9 OF 12 MEDLINE DUPLICATE 3  
 AN 1999328203 MEDLINE  
 DN 99328203  
 TI Randomized trial of sildenafil for the treatment of **erectile** dysfunction in **spinal** cord injury. Sildenafil Study Group.  
 AU Giuliano F; Hultling C; El Masry W S; Smith M D; Osterloh I H; Orr M; Maytom M  
 CS Service d'Urologie, AP-HP, CHU de Bicetre, Le Kremlin Bicetre, France.  
 SO ANNALS OF NEUROLOGY, (1999 Jul) 46 (1) 15-21.  
 Journal code: 6AE. ISSN: 0364-5134.  
 CY United States  
 DT (CLINICAL TRIAL)  
 Journal; Article; (JOURNAL ARTICLE)  
 (RANDOMIZED CONTROLLED TRIAL)  
 LA English  
 FS Priority Journals  
 EM 199910  
 EW 19991001  
 AB **Erectile** dysfunction is a common complication of **spinal** cord injury. This double-blind, placebo-controlled, two-way crossover study assessed the efficacy and safety of oral sildenafil in men with

**erectile** dysfunction caused by traumatic **spinal** cord injury. A total of 178 men (mean age, 38 years) received placebo or sildenafil 1 hour before sexual activity for 6 weeks; after a 2-week washout period, the men received the alternate treatment for 6 weeks. The 50-mg starting dose could be adjusted to 100 or 25 mg based on efficacy and tolerability. Efficacy was assessed by using global efficacy questions, the International Index of **Erectile** Function (IIEF), and a patient log of **erectile** activity. Of 143 men with residual **erectile** function at baseline, 111 (78%) reported improved **erections** and preferred sildenafil to placebo. For all men (including those who reported no residual **erectile** function at baseline), 127 of 168 (76%) reported improved **erections** and preferred sildenafil to placebo. For all men, 132 of 166 (80%) reported that sildenafil improved sexual intercourse compared with 17 of 166 men (10%) reporting improvement with placebo. IIEF questions assessing the ability to achieve and maintain **erections** and satisfaction with sexual intercourse demonstrated significant improvement with sildenafil. Sildenafil was well tolerated, with a low rate of discontinuation because of treatment-related adverse events (2% vs 1% for placebo). Oral sildenafil is an effective and well-tolerated treatment for **erectile** dysfunction caused by **spinal** cord injury.

TI Randomized trial of sildenafil for the treatment of **erectile** dysfunction in **spinal** cord injury. Sildenafil Study Group.

AB **Erectile** dysfunction is a common complication of **spinal** cord injury. This double-blind, placebo-controlled, two-way crossover study assessed the efficacy and safety of oral sildenafil in men with **erectile** dysfunction caused by traumatic **spinal** cord injury. A total of 178 men (mean age, 38 years) received placebo or sildenafil 1 hour before sexual activity. . . . or 25 mg based on efficacy and tolerability. Efficacy was assessed by using global efficacy questions, the International Index of **Erectile** Function (IIEF), and a patient log of **erectile** activity. Of 143 men with residual **erectile** function at baseline, 111 (78%) reported improved **erections** and preferred sildenafil to placebo. For all men (including those who reported no residual **erectile** function at baseline), 127 of 168 (76%) reported improved **erections** and preferred sildenafil to placebo. For all men, 132 of 166 (80%) reported that sildenafil improved sexual intercourse compared with 17 of 166 men (10%) reporting improvement with placebo. IIEF questions assessing the ability to achieve and maintain **erections** and satisfaction with sexual intercourse demonstrated significant improvement with sildenafil. Sildenafil was well tolerated, with a low rate of discontinuation because of treatment-related adverse events (2% vs 1% for placebo). Oral sildenafil is an effective and well-tolerated treatment for **erectile** dysfunction caused by **spinal** cord injury.

CT Check Tags: Human; Male; Support, Non-U.S. Gov't  
Administration, Oral  
Cross-Over Studies  
Double-Blind Method  
\*Impotence: DT, drug therapy  
Impotence: ET, etiology  
Phosphodiesterase Inhibitors: AD, administration & dosage  
Phosphodiesterase Inhibitors: AE, adverse effects  
\*Phosphodiesterase Inhibitors: TU, therapeutic use  
Piperazines: AD, administration & dosage  
Piperazines: AE, adverse effects  
\*Piperazines: TU, therapeutic use  
\*Spinal Cord Injuries: CO, complications

RN 139755-83-2 (sildenafil)

L9 ANSWER 10 OF 12 MEDLINE DUPLICATE 4  
AN 1999071156 MEDLINE  
DN 99071156  
TI Efficacy and safety of oral sildenafil (Viagra) in men with **erectile** dysfunction caused by **spinal** cord injury.

AU Derry F A; Dinsmore W W; Fraser M; Gardner B P; Glass C A; Maytom M C; Smith M D

CS National Spinal Injury Centre, Stoke Mandeville, UK.

SO NEUROLOGY, (1998 Dec) 51 (6) 1629-33.  
Journal code: NZ0. ISSN: 0028-3878.

CY United States

DT (CLINICAL TRIAL)  
Journal; Article; (JOURNAL ARTICLE)  
(RANDOMIZED CONTROLLED TRIAL)

LA English

FS Abridged Index Medicus Journals; Priority Journals

EM 199903

EW 19990301

AB OBJECTIVE: To evaluate the efficacy and safety of 50-mg doses of sildenafil during a 28-day period in patients with **erectile** dysfunction caused by **spinal** cord injury (cord level range, T6 through L5). BACKGROUND: Sildenafil is an orally active, potent, and selective inhibitor of phosphodiesterase type 5, an important regulator of cyclic guanosine monophosphate in the human corpus cavernosum. METHODS: To be included in this double-blind, placebo-controlled study, all patients had to be able to achieve at least a partial reflexogenic **erectile** response to penile vibratory stimulation. The study utilized a single triangular sequential trial design. A total of 27 patients were randomized to receive 50 mg of sildenafil or placebo, taken orally as required (not more than once daily) approximately 1 hour before sexual activity. RESULTS: After 28 days of treatment, nine of 12 patients (75%) on sildenafil and one of 14 patients (7%) on placebo reported that treatment had improved their **erections** (p=0.0043). Furthermore, eight of 12 patients (67%) on sildenafil and two of 13 patients (15%) on placebo indicated that they wished to continue treatment (p=0.018). A significant improvement in satisfaction with their sex life was reported by patients taking sildenafil (p=0.012). No patients discontinued treatment due to adverse events. CONCLUSION: Oral sildenafil, taken as required (not more than once daily), significantly improves the quality of **erections** and satisfaction with sex life in men with **erectile** dysfunction caused by a **spinal** cord injury between T6 and L5.

TI Efficacy and safety of oral sildenafil (Viagra) in men with **erectile** dysfunction caused by **spinal** cord injury.

AB OBJECTIVE: To evaluate the efficacy and safety of 50-mg doses of sildenafil during a 28-day period in patients with **erectile** dysfunction caused by **spinal** cord injury (cord level range, T6 through L5). BACKGROUND: Sildenafil is an orally active, potent, and selective inhibitor of phosphodiesterase. . . be included in this double-blind, placebo-controlled study, all patients had to be able to achieve at least a partial reflexogenic **erectile** response to penile vibratory stimulation. The study utilized a single triangular sequential trial design. A total of 27 patients were. . . of 12 patients (75%) on sildenafil and one of 14 patients (7%) on placebo reported that treatment had improved their **erections** (p=0.0043). Furthermore, eight of 12 patients (67%) on sildenafil and two of 13 patients (15%) on placebo indicated that they. . . due to adverse events. CONCLUSION: Oral sildenafil, taken as required (not more than once daily), significantly improves the quality of **erections** and satisfaction with sex life in men with **erectile** dysfunction caused by a **spinal** cord injury between T6 and L5.

CT Check Tags: Human; Male; Support, Non-U.S. Gov't  
Administration, Oral  
Adult  
Double-Blind Method  
\*Enzyme Inhibitors: AE, adverse effects  
\*Impotence: DT, drug therapy

\*Impotence: ET, etiology  
 Middle Age  
 \*Piperazines: AE, adverse effects  
 Reflex: DE, drug effects  
 Sexuality  
 \*Spinal Cord Injuries: CO, complications  
 RN 139755-83-2 (sildenafil)

L9 ANSWER 11 OF 12 MEDLINE  
 AN 1999024309 MEDLINE  
 DN 99024309  
 TI Oral sildenafil (Viagra) on trial.  
 AU van der Linde I  
 SO SOUTH AFRICAN MEDICAL JOURNAL, (1998 Oct) 88 (10) 1290.  
 Journal code: U4R. ISSN: 0038-2469.  
 CY South Africa  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199903  
 EW 19990303  
 CT Check Tags: Female; Human; Male  
 \*Enzyme Inhibitors: TU, therapeutic use  
 Follow-Up Studies  
 \*Impotence: DT, drug therapy  
 \*Piperazines: TU, therapeutic use  
 Spinal Cord Injuries: CO, complications  
 RN 139755-83-2 (sildenafil)

L9 ANSWER 12 OF 12 HCAPLUS COPYRIGHT 2000 ACS  
 AN 1999:68946 ~~HCAPLUS~~  
 DN 130:231768  
 TI Pharmacotherapy of male **erectile** dysfunction with sildenafil  
 AU Kulkarni, S. K.; Reddy, D. S.  
 CS University Institute of Pharmaceutical Sciences, Panjab University,  
 Chandigarh, 160 014, India  
 SO Indian J. Pharmacol. (1998), 30(6), 367-378  
 CODEN: INJPD2; ISSN: 0253-7613  
 PB Indian Pharmacological Society  
 DT Journal; General Review  
 LA English  
 AB A review with 55 refs. Male **erectile** dysfunction (MED) is a  
 common sexual disorder influenced by psychol., org., phys., endocrine and  
 neurovascular factors. Parasympathetic stimulation activates cholinergic  
 receptors resulting in increased prodn. of nitric oxide and vasoactive  
 peptides which increase the cGMP (cGMP) and thereby increasing vascular  
 smooth muscle relaxation and vasodilation of corpus cavernosum. This  
 causes rigid penile **erection** sufficient for sexual intercourse.  
 However, in **erectile** dysfunction, there is an inability to  
 achieve or maintain a penile **erection**. One of the current  
 approaches in therapies for MED includes elevating levels of cGMP using  
 phosphodiesterase (PDE) inhibitors. Sildenafil is the first orally  
 active, potent and competitive inhibitor of type-5 cGMP-specific PDE  
 enzyme that has been recently approved for the treatment of MED.  
 Sildenafil is a synthetic methylpiperazine deriv. that selectively  
 inhibits PDE in human corpus cavernosum. Sildenafil has demonstrated its  
 effectiveness in **erectile** dysfunction in several preclin. and  
 clin. studies. It is well-tolerated (50-100 mg/day, p.o.) and safe agent  
 for **erectile** dysfunction in patients with diabetes, traumatic  
**spinal cord injury**, psychol. causes and physiol. disorders.  
 Adverse events reported include transient headache, dyspepsia, flushing,  
 diarrhea and visual disturbance. The discovery of sildenafil has not  
 only resulted in a huge market for drugs, but also unfolded the pathophysiol.  
 of **erectile** dysfunction. However, more controlled clin. studies



are needed to establish the safety of sildenafil in patients with different age groups.  
TI Pharmacotherapy of male **erectile** dysfunction with sildenafil  
AB A review with 55 refs. Male **erectile** dysfunction (MED) is a common sexual disorder influenced by psychol., org., phys., endocrine and neurovascular factors. Parasympathetic stimulation activates cholinergic.

. . . increase the cGMP (cGMP) and thereby increasing vascular smooth muscle relaxation and vasodilation of corpus cavernosum. This causes rigid penile **erection** sufficient for sexual intercourse. However, in **erectile** dysfunction, there is an inability to achieve or maintain a penile **erection**. One of the current approaches in therapies for MED includes elevating levels of cGMP using phosphodiesterase (PDE) inhibitors. Sildenafil is. . . Sildenafil is  
a synthetic methylpiperazine deriv. that selectively inhibits PDE in human corpus cavernosum. Sildenafil has demonstrated its effectiveness in **erectile** dysfunction in several preclin. and clin. studies. It is well-tolerated (50-100 mg/day, p.o.) and safe agent for **erectile** dysfunction in patients with diabetes, traumatic **spinal** cord injury, psychol. causes and physiol. disorders. Adverse events reported include transient headache, dyspepsia, flushing, diarrhea and visual disturbance. The discovery of sildenafil has not only resulted in a huge market for drugs, but also unfolded the pathophysiol. of **erectile** dysfunction. However, more controlled clin. studies are needed to establish the safety of sildenafil in patients with different age groups.  
ST review sildenafil **erectile** dysfunction

IT **Impotence**

(sildenafil treatment **erectile** dysfunction in men)

IT 139755-83-2, Sildenafil

RL: ADV (Adverse effect, including toxicity); BAC (Biological activity or effector, except adverse); THU (Therapeutic use); BIOL (Biological study);

USES (Uses)

(sildenafil treatment **erectile** dysfunction in men)

IT 9068-52-4, CGMP Phosphodiesterase

RL: BPR (Biological process); BIOL (Biological study); PROC (Process)  
(sildenafil treatment **erectile** dysfunction in men)

=>

=>

Executing the logoff script...

=> LOG H

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	14.65	22.30
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-1.67	-1.67

SESSION WILL BE HELD FOR 60 MINUTES  
STN INTERNATIONAL SESSION SUSPENDED AT 14:43:36 ON 26 MAR 2000